



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

McCarthy, Alexandra L., Isenring, Elisabeth, Baumann, Sally, Yates, Patsy, Skerman, Helen M., Walpole, Euan, & Thomson, Damien B (2013) Nutrition screening and risk factors prior to chemotherapy in cancer patients 65 years and older. *Supportive Care in Cancer*, 21(S1), S220.

This file was downloaded from: <http://eprints.qut.edu.au/60758/>

© Copyright 2013 Springer

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

MASCC 2013 Abstract

Limits: 250 word text, 20 word title, 1 image or graph (50 words/image).

Authors: McCarthy A¹, Isenring E², Baumann S², Patsy Yates¹, Helen Skerman¹, Euan Walpole³, Damien Thomson³

¹School of Nursing, Queensland University of Technology, Brisbane, Australia, ²Centre for Dietetics Research, School of Human Movement Studies, The University of Queensland, Brisbane, Queensland, Australia, ³ Cancer Care Services, Princess Alexandra Hospital, Metro South, Brisbane, Queensland, Australia.

Topic: Nutrition

Nutrition screening and risk factors prior to chemotherapy in cancer patients 65 years and older

Background

The largest proportion of cancer patients are aged 65 years and over. Increasing age is also associated with nutritional risk and multi-morbidities—factors which complicate the cancer treatment decision-making process in older patients.

Objectives

To determine whether malnutrition risk and Body Mass Index (BMI) are associated with key oncogeriatric variables as potential predictors of chemotherapy outcomes in geriatric oncology patients with solid tumours.

Methods

In this longitudinal study, geriatric oncology patients (aged ≥ 65 years) received a Comprehensive Geriatric Assessment (CGA) for baseline data collection prior to the commencement of chemotherapy treatment. Malnutrition risk was assessed using the Malnutrition Screening Tool (MST) and BMI was calculated using anthropometric data. Nutritional risk was compared with other variables collected as part of standard CGA. Associations were determined by chi-square tests and correlations.

Results

Over half of the 175 geriatric oncology patients were at risk of malnutrition (53.1%) according to MST. BMI ranged from 15.5–50.9kg/m², with 35.4% of the cohort overweight when compared to geriatric cutoffs. Malnutrition risk was more prevalent in those who were underweight (70%) although many overweight participants presented as at risk (34%). Malnutrition risk was associated with a diagnosis of colorectal or lung cancer ($p=0.001$), dependence in activities of daily living ($p=0.015$) and impaired cognition ($p=0.049$). Malnutrition risk was positively associated with vulnerability to intensive cancer therapy ($\rho=0.16$, $p=0.038$). Larger BMI was associated with a greater number of multi-morbidities ($\rho=.27$, $p=0.001$).

Conclusions

Malnutrition risk is prevalent among geriatric patients undergoing chemotherapy, is more common in colorectal and lung cancer diagnoses, is associated with impaired functionality and cognition and negatively influences ability to complete planned intensive chemotherapy.

Key words

Geriatric oncology; malnutrition risk; CGA; nutrition screening; chemotherapy